



## Responder Health Advisory Haiti Earthquake

# Insect Precautions<sup>1</sup>

In the tropics, insects can transmit significant illnesses such as malaria, dengue, yellow fever, and rickettsial disease—some potentially life-threatening. These diseases are best prevented by personal protective measures. In some cases (e.g., malaria or yellow fever), a preventive drug or vaccine is available as well but should never replace personal protection measures. Travelers to areas where insects that transmit these diseases may be present can help minimize their risk by following the insect precautions and protective measures discussed below.

### PERSONAL PROTECTION MEASURES

- Wear clothing that exposes as little skin as is practicable.
- Apply a repellent containing the insecticide DEET (concentration 30 to 35%) or picaridin (concentration 20% or greater for tropical travelers).
  - Picaridin has a pleasant smell, an advantage over DEET.
- The repellent should be applied to all exposed non-sensitive areas of the body. Frequent application ensures continuous protection.
- The time of day and type of insects to be avoided determine when the repellent should be applied.
  - Mosquitoes that transmit malaria (*Anopheles* mosquitoes) are night biters. Thus, if traveling in a malarious area, be especially vigilant in applying repellent from dusk to dawn.
  - Mosquitoes that transmit dengue (*Aedes* mosquitoes) are day biters, and travelers need to be especially vigilant applying repellent during daytime hours when in areas of dengue risk. Peak biting times are usually during the early morning hours and again from late afternoon to dusk.
- Treat outer clothing with permethrin (or other pyrethroid) when traveling in an area of high risk for malaria or other mosquito-borne or tick-borne diseases.
- If you are not sleeping in a sealed, air-conditioned room, sleep under a permethrin-impregnated bed net when at risk of malaria. Regularly check the net for rips and tears, and keep it tucked in around the bed at all times. Ensure that all open windows have insect screens.
- Use an aerosol insecticide before going to bed and a vaporizer device throughout the night.
- Outdoors, a smoldering pyrethroid coil can be used to reduce flying insects.
- In areas where tick-borne disease is a risk, perform a full body check at least once a day.

---

<sup>1</sup> Adapted from information from CDC, Shoreline Medical, and International Travel Medical Society



## INSECTICIDES

The most effective repellents contain **DEET** (N, N diethylmeta-toluamide) or **picaridin** (Cutter Advanced\*, Cutter Advanced Sport\*, KBR3023, Bayrepel, Autan, (RS)-sec-butyl 2-(2-hydroxyethyl). Picaridin is now considered to have comparable efficacy and duration of protection to DEET *at the same concentration*. Both compounds have now been shown to be effective under actual field conditions in tropical countries against both *Anopheles* and *Aedes aegypti* mosquitoes.

- **Duration of protection:**

With both DEET and picaridin, the duration of effectiveness increases as the concentration of repellent increases. With DEET, the effect on duration of protection plateaus at about 50% concentration. Products with less than about 20% picaridin or DEET have a relatively short duration of protection and should be discouraged for use in tropical travelers. The optimal concentration of DEET is considered to be 30-35%. When used by tropical travelers in *appropriate concentrations* (that is, 20% or greater), picaridin should be applied every 4-6 hours. In the U.S., there are products containing 7% picaridin (Cutter Advanced\*) and 15% picaridin (Cutter Advanced Sport\*). The EPA has limited the frequency of application of these 2 products, based on limits for children, as follows: the 7% product should be applied only 3 times per day and the 15% product should be applied only once daily.

- **Use in pregnancy and breastfeeding:**

Picaridin can be used by pregnant women but should not be applied directly to the abdomen. DEET has been shown in 1 study to be safe in the second and third trimesters of pregnancy when used at concentrations of 20% or lower, but the use of DEET in the first trimester has not been well studied. If extensive exposure is anticipated, pregnant women should consider more frequent repellent application since lower concentrations are often used. Neither DEET nor picaridin should be used while breastfeeding, due to a lack of data.

- **Safety:**

DEET is effective against mosquitoes, ticks, fleas, and chiggers and is a remarkably safe insect repellent; only 30 cases of severe toxicity have been reported among billions of uses over 30 years. Most cases of toxic encephalopathy or seizures were reported in young children in whom excessive amounts were used over prolonged periods. There is no long-term information available on the use of picaridin but toxicity tests in animals have shown it to be extremely safe.

The use of another repellent, **IR3535** (3-(N-acetyl-N-butyl) aminopropionic acid ethyl ester; Bug Guard Plus\*), is more controversial, and conflicting data exist over its effectiveness. IR3535 is recommended by WHO as equivalent to picaridin at the same concentration; CDC does not mention or recommend the use of IR3535. IR3535 repellents can be used in children  $\geq 6$  months of age, but due to a lack of data, they should not be used by pregnant or nursing women. In the U.S., IR3535 repellents are available in a range of concentrations up to 20%.

The following precautionary measures can minimize the possibility of adverse reactions to insect repellent containing DEET or picaridin:

- Use repellents according to label directions.
- Apply repellents sparingly and only to exposed skin or clothing.
- Repellents should not be inhaled or ingested and contact with the eyes should be avoided.
- Avoid applying repellents to portions of children's hands that are likely to have contact with eyes or mouth.
- Never use repellents on wounds or irritated skin.
- Wash repellent-treated skin after coming indoors if there is no further risk of exposure to insects.



## DEPARTMENT OF HEALTH & HUMAN SERVICES

---

Federal Occupational Health Service

- If a suspected reaction to insect repellent occurs, wash treated skin and seek medical attention.
- Pregnant and nursing women should minimize use of repellents since about 6-9% of the chemical is absorbed through the skin.

Travelers also should purchase a pyrethroid-containing flying-insect spray to use in living and sleeping areas during evening and nighttime hours.

For added protection against mosquitoes, bednets and clothing may be soaked in or sprayed with permethrin. Permethrin is an insecticide licensed for use on clothing; when applied according to directions it can be effective on clothing for several months and on bednets for half a year. Permethrin physically binds to the fabric, which then can be repeatedly washed without loss of effect; this also prevents absorption through skin. In some countries, deltamethrin liquid is available.

\* Use of brand names is for informational purposes only and does not constitute preference for one brand over another.